
Measuring Uncertainty about Long-Run Predictions

Ulrich K. Müller and Mark W. Watson
Princeton University

Presentation at ECB

June 14, 2014

Set-up


- Observe data x_t , $t = 1, \dots, T$, such as growth rates of GDP, or inflation
- Want to forecast *average* over the next $\lfloor \lambda T \rfloor$ periods

$$f = \lfloor \lambda T \rfloor^{-1} \sum_{l=1}^{\lfloor \lambda T \rfloor} x_{T+l}$$

where $\lambda = 0.5$, say.

- Aim: Construct interval from data $\{x_t\}_{t=1}^T$ that contains f with, say, 90% probability in repeated samples

US Postwar GDP Per Capita



Focus and Challenges

- This paper: statistical (rather than "structural") univariate long-term forecasting
- Econometric challenges
 - Only limited sample information about long-term behavior
 - Set of plausible models of long-term behavior?
 - How to deal with model and/or parameter uncertainty?


Low-Frequency Transformations

- Intuitively, question concerns low-frequency properties of x_t .
- Extract relevant information by computing low-frequency transforms (Müller and Watson, 2008)


$$X_j = T^{-1} \sum_{t=1}^T \sqrt{2} \cos(\pi jt/T) x_t, \quad j = 1, \dots, q$$

where q is a number like $q = 12$, and treat $(X_1, \dots, X_q)'$ and $\hat{\mu} = T^{-1} \sum_{t=1}^T x_t$ as only available data


Cosine Weights




GDP



$q = 12$ LF Transforms for GDP



GDP LF Projection



Pros and Cons of LF Transforms

- Extract low-frequency information in $\{x_t\}$
- Avoids modelling and potential misspecification of higher frequency aspects
- Captures notion that relevant sample information about long-run forecasts limited
- But potential loss of efficiency (see paper)

Standard I(0) Asymptotics for Time Series

- Under a range of primitive conditions on the dependent and heterogeneous mean-zero process $\{u_t\}$, a Central Limit Theorem holds for all fractions of the sample, i.e. for all $0 \leq r_1 < r_2 \leq s_1 < s_2$,

$$\begin{pmatrix} \frac{1}{\sqrt{T}} \sum_{t=\lfloor r_1 T \rfloor + 1}^{\lfloor r_2 T \rfloor} u_t \\ \frac{1}{\sqrt{T}} \sum_{t=\lfloor s_1 T \rfloor + 1}^{\lfloor s_2 T \rfloor} u_t \end{pmatrix} \Rightarrow \mathcal{N} \left(0, \begin{pmatrix} \sigma^2(r_2 - r_1) & 0 \\ 0 & \sigma^2(s_2 - s_1) \end{pmatrix} \right)$$

- This (almost) implies the "Functional" Central Limit Theorem for nicely behaved I(0) processes

$$T^{-1/2} \sum_{t=1}^{\lfloor \cdot T \rfloor} u_t \Rightarrow \sigma W(\cdot)$$

Implications for Low-Frequency Transformations

- Suppose $x_t = \mu + u_t$ and u_t is I(0) in the sense $T^{-1/2} \sum_{t=1}^{\lfloor \cdot T \rfloor} u_t \Rightarrow \sigma W(\cdot)$

- Cosine weights are orthogonal to constant:

$$X_j = \frac{\sqrt{2}}{\sqrt{T}} \sum_{t=1}^T \cos(\pi jt/T) x_t = \frac{\sqrt{2}}{\sqrt{T}} \sum_{t=1}^T \cos(\pi jt/T) u_t$$

- With $\hat{\mu} = T^{-1} \sum_{t=1}^T x_t$, $f = \lfloor \lambda T \rfloor^{-1} \sum_{l=1}^{\lfloor \lambda T \rfloor} x_{T+l}$, $X_0 = \sqrt{T}(\hat{\mu} - \mu)$, and $Y = \sqrt{T}(f - \mu)$, we obtain

$$(X_0, \dots, X_q, Y)' \Rightarrow \mathcal{N}(0, \sigma^2 \Sigma)$$

since weighted averages of Gaussian processes are multivariate Gaussian

- If μ and $\sigma^2 \Sigma$ were known, then could simply report 90% set of the (suitably scaled and centered) conditional normal distribution $Y | \{X_j\}_{j=0}^q$

Invariance

- Impose scale and translation invariance:

$$\{x_t\}_{t=1}^T \mapsto \{m + cx_t\}_{t=1}^T \quad \text{for any } m \text{ and } c \neq 0$$

must lead to corresponding transformation of predictive set

- Can show: Under invariance, asymptotic problem becomes construction of prediction set of

$$Y^s = \frac{Y - X_0}{s_X} \text{ given } X^s = \left(\frac{X_1}{s_X}, \dots, \frac{X_q}{s_X} \right)'$$

where $s_X^2 = q^{-1} \sum_{j=1}^q X_j^2$

⇒ Invariance takes care of lack of knowledge of μ and σ (but still need to know Σ to compute the conditional distribution)

Low-Frequency Forecasts—I(0) Model

- In I(0) model, it turns out that


- $Y^s = \frac{Y - X_0}{s_X}$ is scaled Student- t_q , scaled by $\sqrt{1 + \lambda^{-1}}$

- X^s is independent of Y^s

⇒ intervals for f are of the form $\hat{\mu} \pm$ student-t quantiles multiplied by

$$\frac{(1 + \lambda^{-1})^{1/2} s_X}{\sqrt{T}}$$

GDP 50% and 90% Intervals in I(0) Model



Beyond the I(0) Model

- Natural concern that I(0) model is “too stationary”
- Assume local-level model

$$x_t = \mu + \frac{g}{T} \sum_{s=1}^t \eta_s + \varepsilon_t = \mu + u_t$$


where $\{\varepsilon_t\}$ and $\{\eta_t\}$ are I(0) with identical long-run variance σ^2 , so that $g \geq 0$ measures extent of local mean variability

- Still implies


$$T^{-1/2} \sum_{t=1}^{[\cdot T]} u_t \Rightarrow \sigma G(\cdot) \quad (1)$$

for Gaussian process G , so that $(X_0, \dots, X_q, Y)' \Rightarrow \mathcal{N}(0, \sigma^2 \Sigma)$, where now $\Sigma = \Sigma(g)$


GDP Predictive Densities, LLM, $\lambda = 0.2$




GDP LF Projection



GDP LF Likelihood in LLM



GDP Bayes Predictive Densities



Beyond the Local-Level Model

- Approach generalizes to any model $x_t = \mu + u_t$ that satisfies

$$T^{-\alpha} \sum_{t=1}^{\lfloor \cdot T \rfloor} u_t \Rightarrow \sigma G(\cdot)$$

for some Gaussian process G and α (for example: fractional model).

- Possible to derive predictive set that remains valid for arbitrary G ? No, since Σ then entirely unconstrained.
- Need some regularity of x_t to be able to forecast.
- Consider covariance stationarity of Δx_t (allowing mean growth rate to vary stochastically).

Local-To-Zero Spectrum

- Let $s_T : [-\pi, \pi] \mapsto \mathbb{R}_+$ be a sequence of (pseudo) spectral densities of $\{x_t\}$, and define the *local-to-zero spectrum* $S : \mathbb{R} \mapsto \mathbb{R}$ via

$$S(\omega) = T^{-2\alpha+1} \lim_{T \rightarrow \infty} s_T(\omega/T). \quad (2)$$

for suitable α .

- Under some linear process conditions on Δx_t and (1), we show

$$(X_0, \dots, X_q, Y)' \Rightarrow \mathcal{N}(0, \sigma^2 \Sigma)$$

where Σ is a function of S .

\Rightarrow Long-run forecasting uncertainty is fully determined by (pseudo) spectral shape close to origin.

Local Spectra

- Fractional model:

$$S_d(\omega) \propto |\omega|^{-2d}, d \in (-0.5, 1.5)$$

- Local-to-Unity model (AR(1) with $\rho_T = 1 - c/T$):

$$S_c(\omega) \propto 1/(\omega^2 + c^2)$$





- Local-Level model $x_t = \mu + b\varepsilon_t + \frac{1}{T} \sum_{s=1}^t \eta_s$:

$$S_b(\omega) \propto 1/\omega^2 + b^2$$

⇒ All these local spectra are in bcd-family

$$S_\theta(\omega) \propto \left(\frac{1}{\omega^2 + c^2} \right)^{2d} + b^2, \quad \theta = (b, c, d)$$

Local Log-Spectra



Parameter Uncertainty

- Local spectrum depends on $\theta = (b, c, d)$, which cannot be estimated consistently by fixed number q of cosine transforms.

- Recall that via invariance, (asymptotic) problem is to forecast $Y^s = \frac{Y - X_0}{s_X}$ by $X^s = \left(\frac{X_1}{s_X}, \dots, \frac{X_q}{s_X}\right)'$, where $s_X^2 = q^{-1} \sum_{j=1}^q X_j^2$.

- Let $\Psi(X^s)$ be a predictive interval of level $1 - \alpha$. Determine Ψ^* that minimizes weighted average expected length over θ , subject to coverage constraint for all values of θ :

$$\min_{\Psi} \int w(\theta) E_{\theta}[\text{length}(\Psi(X^s))] d\theta \quad \text{s.t.} \quad P_{\theta}(Y^s \in \Psi(X^s)) \geq 1 - \alpha \quad \forall \theta \in \Theta$$

\Rightarrow Almost same problem as in Müller, Elliott and Watson (2013)

Parameter Uncertainty: Conditional Properties

- Potential problem: $\Psi^*(X^s)$ could be empty for some X^s , and have otherwise unreasonable conditional properties
 - ⇒ generic potential problem of descriptions of uncertainty in nonstandard problems with sets that (only) satisfy confidence type property
 - ⇒ see Müller and Norets (2012)
- Solution: Impose that $\Psi^*(X^s)$ contains the $1 - a$ credible set relative to the prior w .

Implementation


- Set $q = 12$.
- Choose weighting function w uniformly distributed on $d \in [-0.4, 1.4]$ in fractional model
 - \Rightarrow seek to minimize expected length on average with data drawn from fractional model, subject to including the $1 - \alpha$ credible set with that prior and model
- Impose coverage $P_\theta(Y^s \in \Psi(X^s)) \geq 1 - \alpha$ in larger class with local-to-zero spectrum

$$S_\theta(\omega) \propto \left(\frac{1}{\omega^2 + c^2} \right)^{2d} + b^2$$


with $d \in [-0.4, 1.4]$ and b, c arbitrary.

\Rightarrow Frequentist robustification of Bayes credible set


GDP 50% Interval




GDP 90% Intervals




US Postwar PCE Inflation





Inflation 50% Intervals





Inflation 90% Intervals



Labor Productivity 50% Interval



Labor Productivity 90% Interval



Conclusions

- Formalization of uncertainty of statistical long-term predictions
 - Low-frequency transformations to yield robustness.
 - Need regularity. Express regularity via shapes of local-to-zero spectrum.
 - Parameter uncertainty resolved by length minimizing robustification of Bayes credible sets.
- Extension to multivariate problem computationally difficult